## Photo-induced Charge Transfer in Azapyrene-Tetrathiafulvalene Triads

P. Zhou<sup>1</sup>, U. Aschauer<sup>1</sup>, T. Feurer<sup>2</sup>, R. Häner<sup>1</sup>, S. Liu<sup>1</sup>\*

<sup>1</sup>Department of Chemistry, Biochemistry and Pharmaceutical Sciences, University of Bern, Freiestrasse 3, 3012 Bern, Switzerland, <sup>2</sup>Institute of Applied Physics, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

Tetrathiafulvalene (TTF)-based donor-acceptor (D-A) ensembles have attracted a lot of attention due to their unique (opto)electronic properties and potential applications in organic semiconductors, photovoltaics, sensors, switches and molecular electronics.<sup>1-3</sup> To develop high-performance electronic devices, control over multiple charge-transfer (CT) pathways in D-A ensembles is of prime importance. Recently, we have demonstrated chemical and ultrafast optical regulation of distinct photo-induced charge flows within such D-A systems.4,5 As a continuation of our ongoing work, we herein describe redox and optical properties of new D-A ensembles (Chart 1) which were prepared by covalent linkage of two TTF donor units to a central azapyrene acceptor either with or without two tert-butyl groups. A detailed experimental and theoretical study of electronic interactions between D and A units and ICT processes in these triads is presented.



**Chart 1.** Chemical structures of triads 2TTF-pyrene and 2TTF-*t*-butyl-pyrene.

[1] Bergkamp, J. J.; Decurtins, S.; Liu, S. X., Current advances in fused tetrathiafulvalene donoracceptor systems. *Chem. Soc. Rev.* **2015**, *44*, 863-74.

[2] Jana, A.; Ishida, M.; Park, J. S.; Bahring, S.; Jeppesen, J. O.; Sessler, J. L., Tetrathiafulvalene-(TTF-) Derived Oligopyrrolic Macrocycles. *Chem. Rev.* **2017**, *117*, 2641-2710.

[3] Pfattner, R.; Pavlica, E.; Jaggi, M.; Liu, S.-X.; Decurtins, S.; Bratina, G.; Veciana, J.; Mas-Torrent, M.; Rovira, C., Photo-induced intramolecular charge transfer in an ambipolar field-effect transistor based on a  $\pi$ -conjugated donor-acceptor dyad. *J. Mater. Chem. C* **2013**, *1*, 3985-3988.

[4] Zhou, P.; Aschauer, U.; Decurtins, S.; Feurer, T.; Häner, R.; Liu, S. X. Chemical control of photoinduced charge-transfer direction in a tetrathiafulvalene-fused dipyrrolylquinoxaline difluoroborate dyad. *Chem. Commun.* **2020**, *56*, 13421-13424.

[5] Rohwer, E. J.; Geng, Y.; Akbarimoosavi, M.; Daku, L. M. L.; Aleveque, O.; Levillain, E.; Hauser, J.; Cannizzo, A.; Haner, R.; Decurtins, S.; Stanley, R. J.; Feurer, T.; Liu, S. X. Optically Controlled Electron Transfer in a Re(I) Complex. *Chem. Eur. J.* **2021,** *27*, 5399-5403.